VOL. 752, NOS. 1 + 2 JOURNAL OF CHROMATOGRAPHY A 1 NOVEMBER 1996

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

REGULAR PAPERS

Column Liquid Chromatography

Study of retention in reversed-phase liquid chromatography using linear solvation energy relationships. I. The stationary phase
by L.C. Tan and P.W. Carr (Minneapolis, MN, USA) and M.H. Abraham (London, UK) (Received 1 May 1996) . . 1

Scale up in preparative chromatography
by C. Heuer, P. Hugo and G. Mann (Berlin, Germany) and A. Seidel-Morgenstern (Magdeburg, Germany) (Received 11 April 1996) 19

Optimizing preparative separations at high recovery yield
by A. Felinger and G. Guiochon (Oak Ridge, TN, USA) (Received 20 May 1996) 31

Investigation of the behaviour of water on the surface of modified silica using differential scanning calorimetry
by C.A. Fung Kee Fung and M.F. Burke (Tucson, AZ, USA) (Received 15 May 1996) 41

Molded continuous poly(styrene–*co*-divinylbenzene) rod as a separation medium for the very fast separation of polymers. Comparison of the chromatographic properties of the monolithic rod with columns packed with porous and non-porous beads in high-performance liquid chromatography of polystyrenes
by M. Petro, F. Svec and J.M.J. Fréchet (Ithaca, NY, USA) (Received 20 May 1996) 59

Assay for determining the number of reactive groups on gels used in affinity chromatography and its application to the optimisation of the epichlorohydrin and divinylsulfone activation reactions
by J.A. Scoble and R.K. Scopes (Bundoora, Australia) (Received 8 May 1996) 67

Chiral recognition models for the liquid chromatographic resolution of π-acidic racemates on a chiral stationary phase derived from *N*-phenyl-*N*-alkylamide of (*S*)-naproxen
by M.H. Hyun, M.S. Na and J.S. Jin (Pusan, South Korea) (Received 5 March 1996) 77

Study of the acetaldehyde induced polymerisation of flavan-3-ols by liquid chromatography–ion spray mass spectrometry
by H. Fulcrand and T. Doco (Montpellier, France), N.-E. Es-Safi (Rabat, Morocco) and V. Cheynier and M. Moutounet (Montpellier, France) (Received 12 April 1996) 85

Enantioseparation of 3-phenylacetylamino-2,6-piperidinedione and related chiral compounds
by Y. Tang, J.C. Reepmeyer, L.K. Revelle and J.A. Wilson (St. Louis, MO, USA) (Received 10 May 1996) 93

Enantioseparation of a protease inhibitor, indinavir, by subcritical fluid chromatography
by L. Chen, R. Thompson, N. Grinberg, B. Johnson and D. Ellison (Rahway, NJ, USA) (Received 21 May 1996) . . 101

Isolation of monoclonal antibodies from cell containing hybridoma broth using a protein A coated adsorbent in expanded beds
by J. Thömmes, A. Bader, M. Halfar, A. Karau and M.-R. Kula (Jülich, Germany) (Received 24 May 1996) 111

Isocratic high-performance liquid chromatography of coenzyme *A* esters involved in the metabolism of 3*S*-hydroxy-3-methylglutaryl-coenzyme A. Detection of related enzyme activities in *Catharanthus roseus* plant cell cultures
by A. Hermans-Lokkerbol, R. van der Heijden and R. Verpoorte (Leiden, Netherlands) (Received 23 May 1996) . . 123

Resolution of enantiomers of uridine analogs, potential antiviral agents
by S. Levin, M. Sterin and A. Magora (Jerusalem, Israel) and A. Popescu (Lund, Sweden) (Received 15 May 1996) . 131

Evaluation of immunoaffinity chromatography as a replacement for organic solvent clean-up of plant extracts for the determination of triazine herbicides by liquid chromatography
by J.F. Lawrence and C. Ménard (Ottawa, Canada) and M.-C. Hennion, V. Pichon, F. LeGoffic and N. Durand (Paris, France) (Received 13 May 1996) 147

High-performance liquid chromatographic analysis of polydisperse ethoxylated non-ionic surfactants in aqueous samples
by T.C.G. Kibbey, T.P. Yavaraski and K.F. Hayes (Ann Arbor, MI, USA) (Received 1 May 1996) 155

Simultaneous determination of common mono- and divalent cations by ion chromatography with an unmodified silica gel column
by K. Ohta, M. Sando and K. Tanaka (Aichi, Japan) and P.R. Haddad (Hobart, Australia) (Received 9 April 1996) . 167

(Continued overleaf)

Contents (continued)

Gas Chromatography

Homologous series methods for determining hold-up parameters in isothermal gas chromatography
by S. Le Vent (Manchester, UK) (Received 13 May 1996) 173

Multi-channel on-line chromatographic analysis system for quick air–fuel ratio measurement
by F. Akdoğan (Bornova-İzmir, Turkey) (Received 7 May 1996) 183

Unusual behaviour of a new kind of side chain crown ether polysiloxanes used in capillary gas chromatography
by P. Jing (Shaanxi Prov., China), R. Fu, W. Zhou and R. Dai (Beijing, China) and Z. Huang and Y. Chen (Wuhan, China) (Received 14 May 1996) 189

Application of correlation analysis for identification of polychlorinated biphenyls
by A. Kubátová, M. Matucha and J.G.K. Ševčík (Prague, Czech Republic) (Received 20 May 1996) 197

High-resolution gas chromatography–nitrogen–phosphorous detection of saturated volatile aldehydes derivatized with 2-hydrazinobenzothiazole
by E.E. Stashenko (Bucaramanga, Colombia), J.W. Wong (Davis, CA, USA), J.R. Martínez (Setauket, NY, USA), A. Mateus (Bucaramanga, Colombia) and T. Shibamoto (Davis, CA, USA) (Received 3 April 1996) 209

Changes in chemical composition of catalytically hydrogenated orange oil (*Citrus sinensis*)
by E.E. Stashenko, R. Martínez, M.H. Pinzón and J. Ramírez (Bucaramanga, Colombia) (Received 13 May 1996) . 217

Volatile secondary metabolites from *Spilanthes americana* obtained by simultaneous steam distillation–solvent extraction and supercritical fluid extraction
by E.E. Stashenko, M.A. Puertas and M.Y. Combariza (Bucaramanga, Colombia) (Received 13 May 1996) 223

Influence of gas chromatographic parameters on measurement of $^{13}C/^{12}C$ isotope ratios by gas–liquid chromatography–combustion isotope ratio mass spectrometry. I
by W. Meier-Augenstein and P.W. Watt (Dundee, UK) and C.-D. Langhans (Heidelberg, Germany) (Received 13 May 1996) 233

Electrophoresis

Linear free energy relationship analysis of microemulsion electrokinetic chromatographic determination of lipophilicity
by M.H. Abraham (London, UK), C. Treiner (Paris, France), M. Roses and C. Rafols (Barcelona, Spain) and Y. Ishihama (Ibaraki, Japan) (Received 21 May 1996) 243

Isotachophoretic crystallization of ionogenic substances in electrolyte solutions
by O.V. Oshurkova, A.I. Gorshkov and V.P. Nesterov (St. Petersburg, Russia) (Received 14 May 1996) 251

Detection of electrophoretically separated amylase inhibitors in starch–polyacrylamide gels
by A.P. Giri and M.S. Kachole (Aurangabad, India) (Received 21 May 1996) 261

Separation of chiral polychlorinated biphenyls by micellar electrokinetic chromatography using β- and γ-cyclodextrin mixtures in the separation buffer
by M.L. Marina, I. Benito, J.C. Díez-Masa and M.J. González (Madrid, Spain) (Received 19 March 1996) 265

Analysis of basic pharmaceuticals by capillary electrophoresis in coated capillaries and on-line mass spectrometric detection
by D. Belder and D. Stöckigt (Mülheim an der Ruhr, Germany) (Received 23 April 1996) 271

Optical resolution of phenylthiohydantoin-amino acids by capillary electrophoresis and identification of the phenylthiohydantoin-D-amino acid residue of [D-Ala^2]-methionine enkephalin
by Y. Kurosu, K. Murayama, N. Shindo, Y. Shisa and N. Ishioka (Tokyo, Japan) (Received 20 May 1996) 279

SHORT COMMUNICATIONS

Column Liquid Chromatography

pH-Dependent elution behaviour of meleme in high-performance cation-exchange chromatography
by S. Ono, Y. Inoue, N. Watanabe, T. Yoshimura and H. Morita (Toiyama, Japan), A. Shiroishi and M. Takakura (Nei-gun, Japan) and C. Shimasaki (Toiyama, Japan) (Received 13 May 1996) 287

New design for the on-line solid-phase extraction of pesticides using membrane extraction disk material and liquid chromatography in environmental waters
by A. Martín-Esteban, P. Fernández and C. Cámara (Madrid, Spain) (Received 26 March 1996) 291

Ion chromatography methods for the simultaneous determination of mineral anions in plant sap
by P. Masson, G. Hilbert and D. Plenet (Villenave d'Ornon, France) (Received 8 May 1996) 298

AUTHOR INDEX VOL. 752 303